

**SWINE FEED
FENBENDAZOLE TYPE C MEDICATED FEED**

INDICATIONS

For the treatment and control of: **Lungworms:** Adult *Metastrongylus apri*, Adult *Metastrongylus pudendotectus*; **Gastrointestinal worms:** Adult and larvae (L3, L4 stages, liver, lung, intestinal forms) large roundworms (*Ascaris suum*), Adult nodular worms (*Oesophagostomum dentatum*, *O. quadrispinulatum*), Adult small stomach worms (*Hyostromylus rubidus*), Adult and larvae (L2, L3, L4 stages - intestinal mucosal forms) whipworms (*Trichuris suis*); and **Kidney worms:** Adult and larvae *Stephanurus dentatus*.

ACTIVE DRUG INGREDIENT

Fenbendazole ^a	10 – 300 g/ton*
---------------------------	-----------------

*The final printed feed label should list only a single drug concentration.

GUARANTEED ANALYSIS

Crude Protein, not less than	_____ %
Lysine, not less than	_____ %
Crude Fat, not less than	_____ %
Crude Fiber, not more than	_____ %
Calcium, not less than	_____ %
Calcium, not more than	_____ %
Phosphorous, not less than	_____ %
Salt ¹ , not less than	_____ %
Salt ¹ , not more than	_____ %
Sodium ² , not less than	_____ %
Sodium ² , not more than	_____ %
Selenium, not less than	_____ ppm
Zinc, not less than	_____ ppm

¹ If added

² Shall be guaranteed only when total Sodium exceeds that furnished by the maximum salt guarantee.


INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed as the sole ration for 3 to 12 consecutive days at the rate of 0.91 to 6.80 pounds daily per 100 pounds of body weight to provide a total dose of 4.08 milligrams fenbendazole per pound of body weight. Type C medicated swine feeds containing fenbendazole can be fed pelleted or as a meal.

WITHDRAWAL PERIODS



Swine must not be slaughtered for human consumption within 4 days following last treatment with this drug product.

OTHER WARNINGS

Parasite resistance may develop to any dewormer. All dewormers require accurate dosing for best results. Following the use of any dewormer, effectiveness of treatment should be monitored. A decrease of effectiveness over time may indicate the development of resistance to the dewormer administered. The parasite management plan should be adjusted accordingly based on regular monitoring.

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

**Manufactured by
Blue Bird Feed Mills
Any town, Indiana, 12345**

NET WEIGHT ON BAG OR INVOICE

Approved by FDA under NADA # 131-675

Lot No.

Rev. 07/2020

^a Sourced from Safe-Guard[®], NADA # 131-675

Safe-Guard[®] is a registered trademark of Intervet International B.V.