

**Tylosin Medicated Liquid Cattle Feed
(Liver Abscesses)
(tylosin phosphate Type B liquid medicated feed)**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Do Not Feed Undiluted

INDICATIONS FOR USE

For reduction of incidence of liver abscesses caused by *Fusobacterium necrophorum* and *Arcanobacterium pyogenes*.

ACTIVE DRUG INGREDIENT

Tylosin (as tylosin phosphate)^a 90 to 360 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than	_____	%
Non-Protein Nitrogen (NPN) ¹ not more than	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ² , not less than.....	_____	%
Salt ² , not more than.....	_____	%
Sodium ³ , not less than.....	_____	%
Sodium ³ , not more than.....	_____	%
Potassium, not less than	_____	%
Vitamin A ^{2,4} , not less than	_____	I.U./lb
Dry Matter, not less than	_____	60%
Dry Matter, not more than	_____	75%
pH	_____	4.5 to 6.0

¹ When added.

² If added.

³ Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴ Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

MIXING DIRECTIONS

Maintain the pH between 4.5 and 6.0.

Thoroughly mix Tylosin Type B Medicated Feed into one ton of complete cattle feed to obtain the correct concentration in the Type C Medicated Feed (Table 1).

For liquid feeds stored in recirculating tank systems: Recirculate immediately prior to use for not less than 10 minutes, moving not less than 1 percent of the tank contents per minute from the bottom of the tank to the top. Recirculate daily as described even when not used.

For liquid feeds stored in mechanical, air or other agitation-type tank systems: Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used.

Table 1: Mixing Directions for Cattle Medicated Feed

Tylosin Concentration in Type B Feed (g/ton)	Desired Tylosin Concentration in Type C Medicated Feed (g/ton)	
	8	10
	Lb. of Type B per ton of Tylosin Medicated Complete Feed	
90	178	223
180	89	112
360	45	56

CAUTION

Do not use in any liquid feed containing sodium metabisulfite or in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

WARNINGS

Withdrawal Periods

No withdrawal period is required when used according to labeling.

To report adverse effects, access medical information, or obtain additional product information, call 1-800-428-4441.

Approved by FDA under NADA # 012-491

MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb. (kg) on bag or bulk

Lot number: (if applicable) _____

Expiration date: (31 days after the date of manufacture) _____

*The medicated feed label must state a single drug concentration.

^aSourced from Tylan™, NADA # 012-491

Tylan and Elanco are trademarks of Elanco or its affiliates.

Revision Date: May 2021