

**Monensin and Bacitracin Methylenedisalicylate
Medicated Broiler Chicken Feed
(monensin and bacitracin methylenedisalicylate Type C
medicated feed)**

INDICATIONS FOR USE

As an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*; and for increase in rate of weight gain and improved feed efficiency in broiler chickens.

ACTIVE DRUG INGREDIENTS

Monensin^a, USP90 to 110 g/ton*
Bacitracin (as feed grade bacitracin methylenedisalicylate)^b 5 to 25 g/ton*

GUARANTEED ANALYSIS

Crude Protein, min.....	_____	%
Lysine, min.....	_____	%
Methionine, min.....	_____	%
Crude Fat, min.....	_____	%
Crude Fiber, max.....	_____	%
Calcium, min.....	_____	%
Calcium, max.....	_____	%
Phosphorus, min.....	_____	%
Salt ¹ , min.....	_____	%
Salt ¹ , max.....	_____	%
Sodium ² , min.....	_____	%
Sodium ² , max.....	_____	%

¹ If added.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed this complete Type C medicated feed as the sole ration.

CAUTION

Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. In the absence of coccidiosis in broiler chickens the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain. Not for broiler breeder replacement chickens.



WARNINGS



Withdrawal Periods and Residue Warnings

No withdrawal period is required when used according to labeling.

Do not feed to laying chickens.

Do not feed to chickens over 16 weeks of age.

Questions/Comments? To report side effects, access medical information, or obtain additional product information, call Huvepharma, Inc. at 1-877-994-4883. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

Approved by FDA under ANADA # 200-797

MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb. (kg) on bag or bulk

Lot number: (if applicable) _____

Do not use this medicated feed after: (90 days from the date of manufacture) _____

Revision Date: October 2024

^a Sourced from Coxidin[®] (monensin Type A medicated article), ANADA # 200-783

^b Sourced from BMD[®] (bacitracin methylenedisalicylate Type A medicated article); NADA # 046-592

*The final printed medicated feed label must state a single drug concentration.

Coxidin is a registered trademark of Huvepharma EOOD.