

Lubabegron Medicated Cattle Feed

(lubabegron Type C medicated feed)

INDICATIONS FOR USE

For reduction of ammonia gas emissions per pound of live weight and hot carcass weight in growing beef steers and heifers fed in confinement for slaughter during the last 14 to 91 days on feed.

Effectiveness has not been demonstrated when fed for less than 14 days.

Ammonia gas emissions were measured for individual animals or small groups of animals held in environmentally controlled facilities. Based on existing information, reliable predictions of the reduction of ammonia gas emissions cannot be made on a herd, farm, or larger scale.

Increased rate of weight gain, improved feed efficiency, and increased carcass leanness have not been demonstrated with this product.

ACTIVE DRUG INGREDIENT

Lubabegron (as lubabegron fumarate)^a..... 1.25 to 4.54 g/ton*

GUARANTEED ANALYSIS

Crude Protein, Min.....	_____	%
Non-Protein Nitrogen (NPN) ¹ , Max... ..	_____	%
Crude Fat, Min... ..	_____	%
Crude Fiber, Max... ..	_____	%
Calcium, Min... ..	_____	%
Calcium, Max... ..	_____	%
Phosphorus, Min... ..	_____	%
Salt ¹ , Min... ..	_____	%
Salt ¹ , Max... ..	_____	%
Sodium ² , Min... ..	_____	%
Sodium ² , Max... ..	_____	%
Potassium ¹ , Min... ..	_____	%
Vitamin A ¹ , Min... ..	_____	I.U./lb

¹ Guarantee required only when nutrient source is added except when the feed is intended, represented, or serves as a principal source of the nutrient.

² Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed 1.25 to 4.54 g/ton (1.39 to 5 ppm) of complete feed (90% dry matter basis) to provide 13-90 mg lubabegron per head per day to growing beef steers and heifers fed in confinement for slaughter as the sole ration during the last 14 to 91 days on feed. The following table provides examples of feeding rates and amount of drug per ton of Type C medicated feed.

Body Weight	Feed Consumption	Lubabegron in Type C Medicated Feed	
		Consumption in mg/hd/day	g/ton
lb	lb/hd/day		
1000	25	57	4.54
1000	25	35	2.8
1000	25	16	1.25
1200	30	68	4.54
1200	30	42	2.8
1200	30	19	1.25
1400	35	79	4.54
1400	35	49	2.8
1400	35	22	1.25

CAUTION

Not approved for use in breeding animals because safety and effectiveness have not been evaluated in these animals. Do not allow horses or other equines access to feed containing lubabegron. A decrease in dry matter intake may be noticed in some animals.

WARNING



Withdrawal Periods: No withdrawal period is required when used according to labeling.



User Safety Warnings: Not for human use. Keep out of reach of children. The active ingredient in Experior, lubabegron, is a beta-adrenergic agonist/antagonist. Individuals with cardiovascular disease should exercise special caution to avoid exposure. When mixing and handling Experior, use protective clothing, impervious gloves, protective eye wear, and a NIOSH-approved dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water; if wearing contact lenses, rinse eyes first, then remove contact lenses and continue to rinse for 5-20 minutes. If irritation persists, seek medical attention. The safety data sheet contains more detailed occupational safety information.

Questions/Comments? To report side effects, access medical information, or obtain additional product information, call Elanco US Inc. at 1-800-428-4441. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

Lot Number: _____

Approved by FDA under NADA # 141-508

MANUFACTURED BY:
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

^a Sourced from ExperiorTM (lubabegron Type A medicated article).

*The final printed medicated feed label must state a single drug concentration.

Experior, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

Last Revision: July 2024