

**Monensin Medicated Feed for Feedlot Cattle  
Feed Efficiency, Coccidiosis  
(monensin Type B medicated feed)**

**Do Not Feed Undiluted**

**INDICATIONS FOR USE**

For improved feed efficiency; and for the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in growing beef steers and heifers fed in confinement for slaughter.

**ACTIVE DRUG INGREDIENT**

Monensin, USP<sup>a</sup>.....41 to 80,000 g/ton\*

**GUARANTEED ANALYSIS**

Crude Protein, min.....	_____	%
Non-Protein Nitrogen (NPN) <sup>1</sup> , max.....	_____	%
Crude Fat, min.....	_____	%
Crude Fiber, max.....	_____	%
Calcium, min.....	_____	%
Calcium, max.....	_____	%
Phosphorus, min.....	_____	%
Salt <sup>2</sup> , min.....	_____	%
Salt <sup>2</sup> , max.....	_____	%
Sodium <sup>3</sup> , min.....	_____	%
Sodium <sup>3</sup> , max.....	_____	%
Potassium, min.....	_____	%
Vitamin A <sup>24</sup> , min.....	_____	I.U./lb

<sup>1</sup> When added.

<sup>2</sup> If added.

<sup>3</sup> Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>4</sup> Other than precursors of Vitamin A.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

## MIXING DIRECTIONS

Thoroughly mix monensin Type B medicated feed with non-medicated feed to manufacture another concentration of Type B medicated feed or a Type C medicated feed. The Type C medicated feed will have a final concentration of 5 to 40 g/ton monensin for improved feed efficiency or 10 to 40 g/ton monensin for the prevention and control of coccidiosis. No additional improvement in feed efficiency has been shown from feeding monensin at levels greater than 30 g/ton (360 mg monensin per head per day).

When using less than 5 lb of Type B medicated feed per ton to manufacture a medicated feed (Type B or Type C medicated feed), it is recommended to further dilute the Type B medicated feed by thoroughly mixing the required amount of Type B medicated feed in a convenient quantity of non-medicated feed, and then adding the remaining quantity of non-medicated feed.

The following table gives examples of how some Type C medicated feed concentrations can be prepared:

Monensin Concentration in Type B Medicated Feed (g/ton)	Resulting Monensin Concentration in Type C Medicated Feed (g/ton)				
	5	10	20	30	40 <sup>†</sup>
	Lb of Type B Medicated Feed to make 1 ton of Type C Medicated Feed				
41	243.9	487.8	975.6	1,463.4	1,951.2
600	16.7	33.3	66.7	100	133.3
2,000	5	10	20	30	40
8,000	1.25 <sup>^</sup>	2.5 <sup>^</sup>	5	7.5	10
80,000	0.125 <sup>^</sup>	0.25 <sup>^</sup>	0.5 <sup>^</sup>	0.75 <sup>^</sup>	1 <sup>^</sup>

<sup>†</sup> No additional improvement in feed efficiency has been shown from feeding monensin at levels greater than 30 g/ton (360 mg monensin per head per day).

<sup>^</sup> When using less than 5 lb of Type B medicated feed per ton, it is recommended to further dilute the Type B medicated feed as described above.

## CAUTION

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

## WARNINGS

### Withdrawal Periods and Residue Warnings

No withdrawal period is required when used according to labeling.

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**Questions/Comments?** To report side effects, access medical information, or obtain additional product information, call Elanco US Inc. at 1-800-428-4441. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

Approved by FDA under NADA # 095-735

**MANUFACTURED BY**  
BLUE BIRD FEED MILL  
Any town, USA 12345

**Net Weight lb. (kg) on bag or bulk**

Lot number: (if applicable) \_\_\_\_\_

Revision Date: June 2023

<sup>a</sup>Sourced from Rumensin™ (monensin Type A medicated article); NADA # 095-735

\*The final printed medicated feed label must state a single drug concentration.

Rumensin and Elanco are trademarks of Elanco or its affiliates.