

**Monensin Medicated Feed for Feedlot Cattle**  
**Coccidiosis**  
**(monensin Type C medicated feed)**

**INDICATIONS FOR USE**

For the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in growing beef steers and heifers fed in confinement for slaughter.

**ACTIVE DRUG INGREDIENT**

Monensin, USP<sup>a</sup> ..... 10 to 40 g/ton\*

**GUARANTEED ANALYSIS**

Crude Protein, min.....	_____	%
Non-Protein Nitrogen (NPN) <sup>1</sup> , max.....	_____	%
Crude Fat, min.....	_____	%
Crude Fiber, max.....	_____	%
Calcium, min.....	_____	%
Calcium, max.....	_____	%
Phosphorus, min.....	_____	%
Salt <sup>2</sup> , min.....	_____	%
Salt <sup>2</sup> , max.....	_____	%
Sodium <sup>3</sup> , min.....	_____	%
Sodium <sup>3</sup> , max.....	_____	%
Potassium, min.....	_____	%
Vitamin A <sup>24</sup> , min.....	_____	I.U./lb

<sup>1</sup> When added.

<sup>2</sup> If added.

<sup>3</sup> Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>4</sup> Other than precursors of Vitamin A.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**FEEDING DIRECTIONS**

Feed this complete Type C medicated feed containing 10 to 40 g monensin per ton to provide 0.14 to 0.42 mg monensin per pound of body weight per day, depending upon severity of challenge, up to a maximum of 480 mg monensin per head per day.

## CAUTION

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

## WARNINGS

### Withdrawal Periods and Residue Warnings

No withdrawal period is required when used according to labeling.  
A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**Questions/Comments?** To report side effects, access medical information, or obtain additional product information, call Elanco US Inc. at 1-800-428-4441. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

Approved by FDA under NADA # 095-735

**MANUFACTURED BY**  
BLUE BIRD FEED MILL  
Any town, USA 12345

**Net Weight lb. (kg) on bag or bulk**

Lot number: (if applicable) \_\_\_\_\_

Revision Date: June 2023

<sup>a</sup>Sourced from Rumensin™ (monensin Type A medicated article); NADA # 095-735

<sup>\*</sup>The final printed medicated feed label must state a single drug concentration.

Rumensin and Elanco are trademarks of Elanco or its affiliates.