

BLUEBIRD  
TYPE C MEDICATED FEED  
FOR WEANED, GROWING & FINISHING PIGS

For the treatment and control of gastrointestinal roundworms (*Ascaris suum*, adults and fourth-stage larvae; *Ascarops strongylina*, adults; *Hyostrogylus rubidus*, adult and fourth-stage larvae; *Oesophagostomum* spp., adult and fourth-stage larvae), kidneyworms (*Stephanurus dentatus*, adults and fourth-stage larvae), lungworms (*Metastrongylus* spp., adults), threadworms (*Strongyloides ransomi*, adults and somatic larvae, and prevention of transmission of infective larvae to piglets, via colostrum or milk, when fed during gestation), lice (*Haematopinus suis*) and mange mites (*Sarcoptes scabiei* var. *suis*).

For increased rate of weight gain and improved feed efficiency in growing finishing swine.

ACTIVE DRUG INGREDIENTS

Ivermectin.....	1.8 /ton
Bacitracin methylene disalicylate .....	30 g/ton

GUARANTEED ANALYSIS

Protein, not less than	___%
Fat, not less than	___%
Fiber, not more than	___%

INGREDIENTS

Each ingredient must be specifically named (unless stated as such in the guaranteed analysis listing) in accordance with the names and definitions adopted by AAFCO. Collective terms as approved by AAFCO may be used where applicable.

N-141097-A-0000 (CVM Retyped) Approved 2/3/1999

Type C Med. Feed for Weaned, Growing + Finishing Pigs (Ivermectin 1.88 /ton

page 2 of 2

Ivermectin/Bacitracin Methylene Disalicylate Combination  
in Medicated Feed for Swine

Type C Medicated Feed

BMD 30g/ton Page 16

Bluebird Labeling

### FEEDING DIRECTIONS

To provide the recommended level of 30 g bacitracin methylene disalicylate/ton and 1.8 g ivermectin/ton (approximately 100 mcg ivermectin per kg of body weight per day), feed as the only feed for 7 consecutive days. When this feed is consumed, resume feeding bacitracin methylene disalicylate.

**WARNING:** Withdraw 5 days before slaughter.  
Keep this and all drugs out of the reach of children.

**CAUTION:** Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

This product contains IVOME<sup>®</sup> brand ivermectin and has been formulated specifically for use in swine only. This product should not be used for other animal species. Do not permit water runoff from swine production sites to directly enter lakes, streams, or ponds. Do not contaminate water by direct application or by the improper disposal of drug containers.

IVOME<sup>®</sup> is a registered trademark of Merial Limited;  
BMD<sup>®</sup> is a registered trademark of Alpharma, Inc.

Manufactured by

BLUEBIRD FEED COMPANY  
City, State, Zip Code

Net Weight on bag or bulk