**Bacitracin Methylenedisalicylate/Narasin/Nicarbazin - Broiler Feed CNE**

(bacitracin methylenedisalicylate and narasin and nicarbazin Type C medicated feed)

## INDICATIONS FOR USE

For the prevention of coccidiosis in broiler chickens caused by *Eimeria necatrix, E. tenella, E. acervulina, E. brunetti, E. mivati,* and *E. maxima;* and as aid in the control of necrotic enteritis caused or complicated by *Clostridium ssp*. or other organisms susceptible to bacitracin methylenedisalicylate.

## ACTIVE DRUG INGREDIENTS

Narasin. ..... 27 - 45 g/ton
Nicarbazin. ..... 27 - 45 g/ton
Bacitracin (as feed grade bacitracin methylenedisalicylate). ..... 100 - 200 g/ton

## GUARANTEED ANALYSIS

Crude Protein (min)..... _____%
Lysine (min) ..... _____%
Methionine (min) ..... _____%
Crude Fat (min) ..... _____%
Crude Fiber (max)..... _____%
Calcium (min)..... _____%
Calcium (max)..... _____%
Phosphorus (min) ..... _____%
Salt[1] (min). ..... _____%
Salt[1] (max). ..... _____%
Sodium[2] (min)..... _____%
Sodium[2] (max)..... _____%

[1]If added
[2]Shall be guaranteed only when total Sodium exceeds that furnished by the maximum salt guarantee.

## INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of Feed Control Officials (AAFCO).

## FEEDING DIRECTIONS

Feed as the sole ration. To control necrotic enteritis, start this feed at first clinical signs of disease; vary bacitracin dosage based on severity of infection; administer for 5-7 days or as long as clinical signs persist, then reduce bacitracin to prevention level (50 g/ton).

## WITHDRAWAL PERIOD

Withdrawal 5 days before slaughter.

**CAUTION**

Do not allow adult turkeys, horses or other equines access to formulations containing narasin. Ingestion of narasin by these species has been fatal. Nicarbazin medicated broilers may show reduced heat tolerance if exposed to high temperature and high humidity. Provide adequate drinking water and ventilation during these periods. For broiler chickens only. Do not feed to laying chickens. Not for use in feeds for breeder chickens. Consult a poultry diagnostic laboratory or pathologist to determine the diagnosis and advice regarding the optimal level of drug.

**Questions/Comments?**

For technical assistance, contact [name of business] at [business telephone number] or [business web address]. For additional information about reporting side effects or other problems with medicated feeds, contact FDA at 1-888-FDA-VETS or http://www.fda.gov/reportanimalae.

Approved by FDA under NADA # 141-124

Lot Number:_____

**MANUFACTURED BY**
Blue Bird Feed Mill
City, State, Zip

**NET WEIGHT ON BAG OR BULK**

30Apr2025