

Monensin Medicated Goat Feed

Type C Medicated Feed

Feed Only to Goats Maintained in Confinement

For the prevention of coccidiosis caused by *Eimeria crandallis*, *E. christensenii*, and *E. ninakohlyakimovae* in goats maintained in confinement.

ACTIVE DRUG INGREDIENT

Monensin, USP..... 20 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Non-Protein Nitrogen (NPN) ¹ , not more than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ² , not less than.....	_____	%
Salt ² , not more than.....	_____	%
Sodium ³ , not less than.....	_____	%
Sodium ³ , not more than.....	_____	%
Copper ^{2,4} , not less than.....	_____	ppm
Copper ^{2,4} , not more than.....	_____	ppm
Selenium, not less than.....	_____	ppm
Vitamin A ^{2,5} , not less than.....	_____	I.U./lb

¹ When added.

² If added.

³ Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴ If total copper exceeds 20 ppm.

⁵ Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed complete feed (20 g/ton) continuously to goats maintained in confinement.

CAUTION

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result.

WARNING

Do not feed to lactating goats.

Approved by FDA under ANADA # 200-639

MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

Lot No. _____