

Monensin Medicated Feedlot Cattle Feed
Type C Medicated Feed
(monensin Type C medicated feed)

For Cattle Fed in Confinement for Slaughter
(For prevention and control of coccidiosis)

INDICATIONS FOR USE

For cattle fed in confinement for slaughter for prevention and control of coccidiosis due to *Eimeria bovis* and *E. zuernii*.

ACTIVE DRUG INGREDIENTS

Monensin, USP 10 to 40 g/ton*

GUARANTEED ANALYSIS

Crude Protein (Min)	_____ %
NPN ¹ (max).....	_____ %
Crude Fat (min).....	_____ %
Crude Fiber (max).....	_____ %
Calcium (min).....	_____ %
Calcium (max).....	_____ %
Phosphorus (min)	_____ %
Salt ² (min)	_____ %
Salt ² (max).....	_____ %
Sodium ³ (min).....	_____ %
Sodium ³ (max).....	_____ %
Potassium (min).....	_____ %
Vitamin A ^{2,4} (min)	_____ IU/lb

¹When added.

²If added.

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials (AAFCO).

FEEDING DIRECTIONS

Feed a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending upon the severity of challenge, up to maximum of 480 milligrams per head per day.

CAUTION

Do not allow horses or other equines access to feeds containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle feed is safe for use in cattle only.

Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle. Must be thoroughly mixed in feeds before use. Do not feed undiluted Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusal and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

WARNING

A withdrawal period has not been established for this product in preruminating calves.
Do not use in calves to be processed for veal.

QUESTIONS/COMMENTS?

For technical assistance, contact Phibro Animal Health Corporation at 888-403-0074 or www.pahc.com. For additional information about reporting side effects or other problems with medicated feeds, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

Approved by FDA under ANADA # 200-559

Lot Number: _____

MANUFACTURED BY

BLUE BIRD FEED MILL
City, State, Zip

Net Weight lb (kg) on bag or bulk

*Final printed label on formulated Type C medicated feed must bear a single drug concentration.

03Sep2025