

Date of Approval: February 9, 2023

# FREEDOM OF INFORMATION SUMMARY

## SUPPLEMENTAL ABBREVIATED NEW ANIMAL DRUG APPLICATION

ANADA 200-701

PARASEDGE™ Multi for Cats  
(imidacloprid + moxidectin)

Topical Solution

Ferrets

This supplement provides for the addition of the following indications for ferrets:  
PARASEDGE™ Multi for Cats is indicated for the prevention of heartworm disease in ferrets caused by *Dirofilaria immitis*. PARASEDGE™ Multi for Cats kills adult fleas (*Ctenocephalides felis*) and is indicated for the treatment of flea infestations on ferrets.

Sponsored by:

Chanelle Pharmaceuticals Manufacturing Ltd.

Table of Contents

I. GENERAL INFORMATION .....	3
II. BIOEQUIVALENCE .....	4
III. HUMAN FOOD SAFETY .....	5
IV. USER SAFETY .....	5
V. AGENCY CONCLUSIONS .....	5

**I. GENERAL INFORMATION**

**A. File Number**

ANADA 200-701

**B. Sponsor**

Chanelle Pharmaceuticals Manufacturing Ltd.  
Loughrea, County Galway, Ireland

Drug Labeler Code: 061651

U.S. Agent Name and Address:  
Dr. James H. Schafer  
Schafer Veterinary Consultants, LLC  
800 Helena Court  
Fort Collins, CO 80524

**C. Proprietary Name**

PARASEDGE™ Multi for Cats

**D. Drug Product Established Name**

imidacloprid + moxidectin

**E. Pharmacological Category**

Antiparasitic

**F. Dosage Form**

Topical solution

**G. Amount of Active Ingredient**

100 mg/mL (10%) imidacloprid and 10 mg/mL (1%) moxidectin

**H. How Supplied**

0.4 mL applicator tube; three tubes per package

**I. Dispensing Status**

Prescription (Rx)

**J. Dosage Regimen**

The recommended minimum dose for a ferret is 9 mg/lb (20 mg/kg) imidacloprid and 0.9 mg/lb (2 mg/kg) moxidectin, once a month, by topical administration.

<b>Ferret (lbs.)</b>	<b>PARASEDGE™ Multi for Cats</b>	<b>Volume (mL)</b>	<b>Imidacloprid (mg)</b>	<b>Moxidectin (mg)</b>
2.0 – 4.4	PARASEDGE™ Multi 9	0.4	40	4

**Only the 0.4 mL applicator tube volume (PARASEDGE™ Multi 9) should be used on ferrets.**

**K. Route of Administration**

Topical

**L. Species/Class**

Ferrets

**M. Indications**

PARASEDGE™ Multi for Cats is indicated for the prevention of heartworm disease in ferrets caused by *Dirofilaria immitis*. PARASEDGE™ Multi for Cats kills adult fleas (*Ctenocephalides felis*) and is indicated for the treatment of flea infestations on ferrets.

**N. Reference Listed New Animal Drug (RLNAD)**

advantage multi™ for cats; imidacloprid + moxidectin; NADA 141-254; Elanco US Inc.

**O. Effect of Supplement**

This supplement provides for the addition of the following indications for ferrets: PARASEDGE™ Multi for Cats is indicated for the prevention of heartworm disease in ferrets caused by *Dirofilaria immitis*. PARASEDGE™ Multi for Cats kills adult fleas (*Ctenocephalides felis*) and is indicated for the treatment of flea infestations on ferrets.

**II. BIOEQUIVALENCE**

CVM did not require additional bioequivalence information for this supplemental approval. The FOI Summary for the original approval of ANADA 200-701, dated June 10, 2021, contains a summary of data that demonstrates bioequivalence of the drug for ferrets.

### **III. HUMAN FOOD SAFETY**

This drug is intended for use in ferrets. Because this new animal drug is not intended for use in food-producing animals, CVM did not require data pertaining to drug residues in food (i.e., human food safety) for approval of this ANADA.

### **IV. USER SAFETY**

The product labeling contains the following information regarding safety to humans handling, administering, or exposed to PARASEDGE™ Multi for Cats:

#### **HUMAN WARNINGS:**

**Not for human use. Keep out of the reach of children.**

**Children should not come in contact with the application site for 30 minutes after application.**

Causes eye irritation. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Exposure to the product has been reported to cause headache; dizziness; and redness, burning, tingling, or numbness of the skin.

**Wash hands thoroughly with soap and warm water after handling.**

If contact with eyes occurs, hold eyelids open and flush with copious amounts of water for 15 minutes. If eye irritation develops or persists, contact a physician. If swallowed, call poison control center or physician immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or physician. People with known hypersensitivity to benzyl alcohol, imidacloprid or moxidectin should administer the product with caution. In case of allergic reaction, contact a physician. If contact with skin or clothing occurs, take off contaminated clothing. Wash skin immediately with plenty of soap and water. Call a poison control center or physician for treatment advice.

The Safety Data Sheet (SDS) provides more detailed occupational safety information. To report suspected adverse drug events, for technical assistance or to obtain a copy of the SDS, contact Virbac AH, Inc. at 1-800-338-3659 or [us.virbac.com](http://us.virbac.com). For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

### **V. AGENCY CONCLUSIONS**

The information submitted in support of this supplemental ANADA satisfy the requirements of section 512(c)(2) of the Federal Food, Drug, and Cosmetic Act. The data demonstrate that PARASEDGE™ Multi for Cats, when used according to the label, is safe and effective for the indications listed in Section I.M. above.