

FREEDOM OF INFORMATION (FOI) SUMMARY

ORIGINAL ABBREVIATED NEW ANIMAL DRUG APPLICATION (ANADA)

ANADA 200-811

MoxiCloprid™ for Cats
(imidacloprid + moxidectin)

Topical Solution

Cats and Ferrets

For Cats: MoxiCloprid™ for Cats is indicated for the prevention of heartworm disease caused by *Dirofilaria immitis*. MoxiCloprid™ for Cats kills adult fleas (*Ctenocephalides felis*) and is indicated for the treatment of flea infestations. MoxiCloprid™ for Cats is also indicated for the treatment and control of ear mite (*Otodectes cynotis*) infestations and the following intestinal parasites:

Intestinal Parasite		Intestinal Stage		
		Adult	Immature Adult	Fourth Stage Larvae
Hookworm Species	<i>Ancylostoma tubaeforme</i>	X	X	X
Roundworm Species	<i>Toxocara cati</i>	X		X

For Ferrets: MoxiCloprid™ for Cats is indicated for the prevention of heartworm disease in ferrets caused by *Dirofilaria immitis*. MoxiCloprid™ for Cats kills adult fleas (*Ctenocephalides felis*) and is indicated for the treatment of flea infestations on ferrets.

Sponsored by:

Bimeda Animal Health Ltd.

Table of Contents

I. GENERAL INFORMATION.....	3
II. BIOEQUIVALENCE	5
III. HUMAN FOOD SAFETY.....	5
IV. USER SAFETY	5
V. AGENCY CONCLUSIONS.....	6

I. GENERAL INFORMATION

A. File Number

ANADA 200-811

B. Sponsor

Bimeda Animal Health Ltd.
1B The Herbert Building
The Park, Carrickmines
Dublin 18, Ireland

Drug Labeler Code: 061133

U.S. Agent Name and Address:

Stephanie Batliner
Bimeda Inc.
One Tower Lane
Suite 2250
Oakbrook Terrace, IL 60181

C. Proprietary Name

MoxiCloprid™ for Cats

D. Drug Product Established Name

imidacloprid + moxidectin

E. Pharmacological Category

Antiparasitic

F. Dosage Form

Topical solution

G. Amount of Active Ingredient

100 mg/mL (10%) imidacloprid and 10 mg/mL (1%) moxidectin

H. How Supplied

0.23 mL applicator tubes; three tubes per carton, 0.4 mL and 0.8 mL applicator tubes;
six tubes per carton

I. Dispensing Status

Prescription (Rx)

J. Dosage Regimen

For cats: The recommended minimum dose for a cat is 4.5 mg/lb (10.0 mg/kg) imidacloprid and 0.45 mg/lb (1.0 mg/kg) moxidectin, once a month, by topical administration.

Cat (lbs.)	Volume (mL)	Imidacloprid (mg)	Moxidectin (mg)
2-5	0.23	23	2.3
5.1-9	0.4	40	4
9.1-18*	0.8	80	8

*Cats over 18 lbs. should be treated with the appropriate combination of MoxiCloprid™ for Cats tubes.

For ferrets: The recommended minimum dose for a ferret is 9 mg/lb (20 mg/kg) imidacloprid and 0.9 mg/lb (2 mg/kg) moxidectin, once a month, by topical administration.

Ferret (lbs.)	Volume (mL)	Imidacloprid (mg)	Moxidectin (mg)
2.0-4.4	0.4	40	4

Only the 0.4 mL applicator tube volume should be used on ferrets.

K. Route of Administration

Topical

L. Species

Cats and ferrets

M. Indications

For cats: MoxiCloprid™ for Cats is indicated for the prevention of heartworm disease caused by *Dirofilaria immitis*. MoxiCloprid™ for Cats kills adult fleas (*Ctenocephalides felis*) and is indicated for the treatment of flea infestations. MoxiCloprid™ for Cats is also indicated for the treatment and control of ear mite (*Otodectes cynotis*) infestations and the following intestinal parasites:

Intestinal Parasite		Intestinal Stage		
		Adult	Immature Adult	Fourth Stage Larvae
Hookworm Species	<i>Ancylostoma tubaeforme</i>	X	X	X
Roundworm Species	<i>Toxocara cati</i>	X		X

For ferrets: MoxiCloprid™ for Cats is indicated for the prevention of heartworm disease in ferrets caused by *Dirofilaria immitis*. MoxiCloprid™ for Cats kills adult fleas (*Ctenocephalides felis*) and is indicated for the treatment of flea infestations on ferrets.

N. Reference Listed New Animal Drug (RLNAD)

advantage multi™ for cats; imidacloprid + moxidectin; NADA 141-254; Elanco US Inc.

II. BIOEQUIVALENCE

The Federal Food, Drug, and Cosmetic Act (FD&C Act), as amended by the Generic Animal Drug and Patent Term Restoration Act (GADPTRA) of 1988, allows for an abbreviated new animal drug application (ANADA) to be submitted for a generic version of an approved new animal drug (RLNAD). The ANADA sponsor is required to show that the generic product is bioequivalent to the RLNAD, which has been shown to be safe and effective. Effectiveness, target animal safety and human food safety data (other than tissue residue data) are not required for approval of an ANADA. If bioequivalence is demonstrated through a clinical endpoint study in a food-producing animal, then a tissue residue study to establish the withdrawal period for the generic product is also required. For certain dosage forms, the agency will grant a waiver from the requirement to perform *in vivo* bioequivalence studies (biowaiver) (55 FR 24645, June 18, 1990; Fifth GADPTRA Policy Letter; Bioequivalence Guideline, October 9, 2002).

Based on the formulation characteristics of the generic product, Bimeda Animal Health Ltd., was granted a biowaiver for the generic product MoxiCloprid™ for Cats (imidacloprid + moxidectin) topical solution. The generic drug product is a topical solution, contains the same active ingredient in the same concentration and dosage form as the RLNAD, and contains no inactive ingredients that may significantly affect the bioavailability of the active ingredient. The RLNAD is advantage multi™ for cats (imidacloprid + moxidectin) topical solution, sponsored by Elanco US Inc., under NADA 141-254, and was approved for use in cats on January 19, 2007, and for use in ferrets on October 31, 2013.

III. HUMAN FOOD SAFETY

This drug is intended for use in cats and ferrets. Because this new animal drug is not intended for use in food-producing animals, CVM did not require data pertaining to drug residues in food (i.e., human food safety) for approval of this ANADA.

IV. USER SAFETY

The product labeling contains the following information regarding safety to humans handling, administering, or exposed to MoxiCloprid™ for Cats:

**Not for human use. Keep out of the reach of children.
Children should not come in contact with the application site for 30 minutes after application.**

Causes eye irritation. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Exposure to the product has been reported to cause headache; dizziness; and redness, burning, tingling, or numbness of the skin. **Wash hands thoroughly with soap and warm water after handling.** If contact with eyes occurs, hold

eyelids open and flush with copious amounts of water for 15 minutes. If eye irritation develops or persists, contact a physician. If swallowed, call poison control center or physician immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or physician. People with known hypersensitivity to benzyl alcohol, imidacloprid or moxidectin should administer the product with caution. In case of allergic reaction, contact a physician. If contact with skin or clothing occurs, take off contaminated clothing. Wash skin immediately with plenty of soap and water. Call a poison control center or physician for treatment advice.

In humans, ocular and dermal irritation, nausea, numbness or tingling of the mouth and lips, anaphylaxis, pruritus, vomiting, and tongue/taste abnormalities have been reported following exposure to this product.

V. AGENCY CONCLUSIONS

The data submitted in support of this ANADA satisfy the requirements of section 512(c)(2) of the FD&C Act. The data demonstrate that MoxiCloprid™ for Cats, when used according to the label, is safe and effective for the conditions of use in the General Information Section above.