


Boehringer Ingelheim AMT ING		PPM Code: L6528
		SKU: 168439
Version: 16	Date: 14.06.2023	Colors
Product:	Senvelgo 1,5mg/ml OS for cats	Black
Component Type/Size:	Leaflet	T
Country(s):	USA	
Dimensions FLAT (HxWxD):	528 x 216 mm	
Dimensions FOLDED:	27 x 176 mm	
Scale/Font Size:	1:1 7.6 pt.	
Spec/Dieline #:	TD2002.03	Finishing/Tech Info
Manufacturer:	CMO Argenta Dundee	
Cancel and Replace:		
Printer Notes:		

Client Information Sheet



Senvelgo[®]

(velagliflozin oral solution)

15 mg/mL

For oral use in cats only

Sodium-glucose cotransporter 2 (SGLT2) inhibitor

The Client Information Sheet contains important information about SENVELGO[®] (velagliflozin oral solution). You should read this information before you start giving SENVELGO to your cat and review it each time the prescription is refilled as there may be new information. This sheet is provided only as a summary and does not take the place of instructions from your veterinarian. Talk with your veterinarian if you do not understand any of this information or if you want to know more about SENVELGO.

What is SENVELGO?

SEVELGO is a drug used to lower high blood glucose (sugar) in otherwise healthy cats with diabetes mellitus not previously treated with insulin. SENVELGO causes sugar to be passed in the urine by preventing the kidneys from pulling sugar back into the blood.

What is diabetes mellitus (commonly called diabetes)?

Diabetes happens when a cat does not make enough insulin (insulin dependent), or the cat's cells do not react correctly to insulin (insulin resistant). Some cats may begin as insulin resistant but may slowly become insulin dependent over time. Sugar is an important source of energy for the cells in your cat's body. Cats with diabetes have high blood sugar (hyperglycemia) because the sugar cannot get into the cells to be used for energy. Insulin helps sugar get into your cat's cells for energy. There is no test available for your veterinarian to know if your cat makes enough insulin or not. Cats with diabetes may show an increased thirst, urination, and hunger; weight loss; and/or weakness in the back legs. **Cats who are insulin dependent should not be treated with SENVELGO because it can cause serious, sometimes fatal, side effects.**

What are some of the possible side effects of SENVELGO?

SEVELGO may be associated with side effects. Serious side effects have happened with or without warning and sometimes led to death in cats treated with SENVELGO. Serious side effects may require your cat to be hospitalized and to be given specific treatments, including insulin. The most common side effects are diarrhea or loose stool, vomiting, and weight loss.

STOP SENVELGO AND CONTACT YOUR VETERINARIAN IMMEDIATELY IF YOU NOTICE ANY OF THE FOLLOWING CHANGES OR SIDE EFFECTS IN YOUR CAT:

- Less interest in food or not eating
- Lack of energy or change in normal activity
- Vomiting or diarrhea
- Weakness, difficulty walking or standing

What should I discuss with my veterinarian before giving SENVELGO?

Tell your veterinarian if:

- Your cat is currently or was previously treated with insulin or other anti-diabetic medications.
- Your cat has been previously diagnosed with diabetic ketoacidosis (a buildup of acids in the blood) or liver or kidney disease.
- Your cat has any other serious health conditions such as acute vomiting, diarrhea, decreased appetite, or weight loss.
- Your cat is taking other medications including prescription drugs and over-the-counter medication.
- Your cat is pregnant, nursing, or you intend to breed him/her.
- Your cat has any planned procedures that involve anesthesia, such as surgery or dentistry.

When should I take my cat to the veterinarian?

- It is very important to take your cat to the veterinarian twice during the first week of taking SENVELGO to monitor your cat for side effects even if your cat seems fine. Your cat should be examined by your veterinarian 2 or 3 days after starting SENVELGO and again about 7 days after starting SENVELGO.
- It is also important to take your cat to the veterinarian immediately if your cat is showing signs of illness including decreased appetite or decreased thirst, vomiting, weakness, decreased activity or depressed attitude.
- It is important to discuss your cat's response to SENVELGO at regular checkups.
- Your veterinarian will determine if your cat is responding as expected and whether your cat should continue using SENVELGO.

What kind of results can I expect when my cat is on SENVELGO for diabetes mellitus?

You may see a decrease in the amount of thirst, urine, and hunger in your cat.

How do I give SENVELGO to my cat?

SEVELGO should be given directly into your cat's mouth with the provided dosing syringe or placed on a small amount of wet food your cat will eat completely (approximately 1 to 3 teaspoons of food). The dosing syringe fits into the bottle opening and has a body weight scale to use for measuring the dose. The body weight scale on the syringe is made to give the amount of SENVELGO prescribed by your veterinarian.

Follow your veterinarian's instructions for how much and how often to give SENVELGO.



- Remove the cap from the bottle and gently push the end of the dosing syringe into the bottle opening.
- Turn the bottle/syringe upside down.
- Pull the plunger back and fill the dosing syringe to the dose prescribed by your veterinarian.
- Turn the bottle upright and remove the dosing syringe from the bottle.



- Put the end of the dosing syringe into your cat's mouth and push the plunger to give the dose.



- Or put the dose next to or on top of a small amount of wet food. Do not mix into food. If SENVELGO is put onto food, make sure your cat eats the food and the entire dose.

If your cat vomits within 30 minutes of giving the SENVELGO dose, your cat can be given the same dose again. If your cat vomits more than 30 minutes after giving SENVELGO, do not give SENVELGO again until the next scheduled dose. After dosing, replace the cap on the bottle and close tightly. If needed, the syringe can be cleaned with a clean, dry cloth.

What should I do if my cat gets more than the prescribed dose?

Contact your veterinarian if you see any side effects. If no side effects are seen, return to the dose prescribed by your veterinarian the following day.

What should I do if my cat gets less than the prescribed dose, or I miss a dose?

- If your cat gets less than the prescribed dose, return to the dose prescribed by your veterinarian the following day.
- If a dose is missed, it should be given as soon as possible on the same day.

How should I store SENVELGO?

SEVELGO can be stored at or below 77°F (25°C) with excursions permitted up to 104°F (40°C). Once the bottle is opened, use the contents within six months.

Keep SENVELGO in a secure location out of reach of dogs, cats, and other animals to avoid accidental ingestion or overdose.

Approved by FDA under NADA # 141-568

SEVELGO[®] is a registered trademark of Boehringer Ingelheim Vetmedica GmbH, used under license.

© 2023 Boehringer Ingelheim Animal Health USA Inc. All rights reserved.

Revised 06/2023

See reverse side for Prescribing Information